



URINALYSIS CONTROL - LEVEL I (URNAL CONTROL I)

CAT. NO. UC5033 **LOT NO.** 1276UC **SIZE:** 12 x 12 ml **EXPIRY:** 2024-05-28

GTIN: 05055273207583

INTENDED USE

This product is intended for in vitro diagnostic use, in the quality control of urine test strips.

DEVICE DESCRIPTION

The Urinalysis Controls are supplied at 2 levels, level 1 and level 2. Ranges are supplied for the following analytes: Albumin, Bilirubin, Blood, Creatinine, Glucose, HCG, Ketones, Leukocytes, Nitrite, pH, Protein, Specific Gravity and Urobilinogen.

SAFETY PRECAUTIONS AND WARNINGS

For *in vitro* diagnostic use only. Do not pipette by mouth. Exercise the normal precautions required for handling laboratory reagents.

Human source material, from which this product has been derived, has been tested at donor level for the Human Immunodeficiency Virus (HIV I, HIV 2) antibody, Hepatitis B Surface Antigen (HbsAg), and Hepatitis C Virus (HCV) antibody and found to be NON-REACTIVE. FDA approved methods have been used to conduct these tests. However, since no method can offer complete assurance as to the absence of infectious agents, this material and all patient samples should be handled as though capable of transmitting infectious diseases and disposed of accordingly.

Health and Safety Data Sheets are available on request.

STORAGE AND STABILITY

OPENED: Store refrigerated (\pm 2°C to \pm 8°C). The Urinalysis Control is stable for 30 days at \pm 2°C to \pm 25°C, or 20 immersions if kept capped in original container and free from contamination. Do not freeze.

UNOPENED: Store refrigerated (+2°C to +8°C). Stable to expiration date printed on individual vials.

PREPARATION FOR USE

The Urinalysis Control is supplied ready for use.

MATERIALS PROVIDED

Urinalysis Control - Level I 12 x 12 ml

ASSIGNED VALUES

Due to the variation caused by test equipment, test reagents and laboratory technique, the quoted ranges are provided for guidance. It is recommended that these ranges are used until each laboratory has established its own ranges, based on individual laboratory requirements.

Each batch of Urinalysis Control is submitted to a number of external laboratories and values are assigned from a consensus of results obtained by these laboratories. With each batch, a control range is provided for individual parameters and each parameter method.





| MANUFACTURER / METHOD | ANALYTE | RANGE |
|---------------------------------|-------------------------------|---|
| 77 Elektronika kft | Bilirubin | NEGATIVE |
| | Blood | NEGATIVE |
| | Ketones | NEGATIVE |
| | Leukocytes | NEGATIVE |
| | Nitrite | NEGATIVE |
| | pΗ | <5 - 6 |
| | Protein | NEGATIVE |
| | Specific Gravity | I.02-I.03 RATIO |
| | Urobilinogen | NORMAL |
| Abbott IMx (intact hCG) | hCG | NEGATIVE |
| 7 tobott ii ix (iiitact ii e a) | lied | THE GIVITY E |
| Abon hCG One Step Ultra | hCG | NEGATIVE |
| | | |
| Acon Mission | Bilirubin | NEGATIVE |
| | Blood | NEGATIVE |
| | Glucose | NEGATIVE |
| | Ketones | NEGATIVE |
| | Leukocytes Nitrite | NEGATIVE NEGATIVE |
| | pH | <5 – 6.5 |
| | Protein | NEGATIVE |
| | Specific Gravity | 1.02-1.03 RATIO |
| | Urobilinogen | NEG – 0.2 mg/dl / NEG – 3.5 μmol/l / NEG – 0.2 EU/dl |
| | - Cr Commogen | 1 1 1 2 1 1 1 1 1 1 2 2 3 3 4 1 1 1 1 2 2 3 3 4 1 1 1 1 1 2 3 3 4 1 1 1 1 1 2 3 3 4 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 |
| Analyticon Combiscreen | Bilirubin | NEGATIVE |
| | Blood | NEGATIVE |
| | Ketones | NEGATIVE |
| | Leukocytes | NEGATIVE |
| | Nitrite | NEGATIVE |
| | pH | <5 - 6 |
| | Protein | NEGATIVE I.02-I.03 RATIO |
| | Specific Gravity Urobilinogen | NORMAL |
| | Orobilinogen | NORMAL |
| Arkray / Menarini | Bilirubin | NEGATIVE |
| 7 a Ki ay 7 i Tenariii | Blood | NEGATIVE |
| | Ketones | NEGATIVE |
| | Leukocytes | NEGATIVE |
| | Nitrite | NEGATIVE |
| | рН | <5 - 6 |
| | Protein | NEGATIVE |
| | Specific Gravity | 1.02 – 1.03 RATIO |
| | Urobilinogen | NORMAL (0.2 – 1 mg/dl) (3.2 – 16 μmol/l) |
| | Orobilinogen | (0.2 - I EU/dl) |
| | | |
| DFI - Cybow | Bilirubin | NEGATIVE |
| | Blood | NEGATIVE |
| | Glucose | NEGATIVE |
| | Ketones | NEGATIVE |
| | Leukocytes | NEGATIVE |
| | Nitrite | NEGATIVE |
| | pH | <5 - 6 |
| | Protein | NEGATIVE |
| | Specific Gravity | 1.02 – 1.03 RATIO |
| | Urobilinogen | NORMAL (0.1 – 1 mg/dl) (1.6 – 16) μmol/l) (0.1 – 1 EU/dl) |
| | _1 | (v.1 – 1 LO/ul) |

| MANUFACTURER / METHOD | ANALYTE | RANGE |
|-----------------------|-----------|----------|
| Dirui H Series | Bilirubin | NEGATIVE |





| | DI ' | NIFC ATIVE |
|------------------------------------|------------------|--------------------------------|
| | Blood | NEGATIVE |
| | Glucose | NEGATIVE |
| | Ketones | NEGATIVE |
| | Leukocytes | NEGATIVE |
| | Nitrite | NEGATIVE |
| | pΗ | <5 - 6.5 |
| | Protein | NEGATIVE |
| | Specific Gravity | 1.02 – 1.03 RATIO |
| | Urobilinogen | NORMAL / +/- / 3.4 – 17 μmol/l |
| EKF Diagnostics QuPID / QuPID Plus | hCG | NEGATIVE |
| | | |
| Erba Lachema | Bilirubin | NEGATIVE |
| | Blood | NEGATIVE |
| | Glucose | NEGATIVE |
| | Ketones | NEGATIVE |
| | Leukocytes | NEGATIVE |
| | Nitrite | NEGATIVE |
| | pН | <5 – 6 |
| | Protein | NEGATIVE |
| | Specific Gravity | 1.02 – 1.03 RATIO |
| | Urobilinogen | NORMAL |
| IND Diagnostic hCG | hCG | NEGATIVE |
| Iris Velocity | Bilirubin | NEGATIVE |
| | Blood | NEGATIVE |
| | Glucose | NEGATIVE |
| | Ketones | NEGATIVE |
| | Leukocytes | NEGATIVE |
| | Nitrite | NEGATIVE |
| | рН | <5-6 |
| | Protein | NEGATIVE |
| | Specific Gravity | 1.015 – 1.025 RATIO |
| | Urobilinogen | NORMAL |
| Min duni | Bilirubin | NEGATIVE |
| Mindray | Blood | NEGATIVE |
| | | NEGATIVE |
| | Glucose | |
| | Ketones | NEGATIVE |
| | Leukocytes | NEGATIVE |
| | Nitrite | NEGATIVE |
| | pH | <5 - 6 |
| | Protein | NEGATIVE LOS DATIO |
| | Specific Gravity | 1.02-1.03 RATIO |
| | Urobilinogen | NORMAL / 0.2 mg / 3.2 μmol/l |
| Nova Century Scientific | hCG | NEGATIVE |
| | + | |

| MANUFACTURER / METHOD | ANALYTE | RANGE |
|--------------------------------|-----------|----------|
| Roche Cobas U601 / Urisys 2400 | Bilirubin | NEGATIVE |
| | Blood | NEGATIVE |
| | Ketones | NEGATIVE |





| | Leukocytes | NEGATIVE |
|----------------------------|------------------|------------------------------------|
| | Nitrite | NEGATIVE |
| | pH | <5 - 6 |
| | Protein | NEGATIVE |
| | | |
| | Specific Gravity | 1.015-1.03 RATIO |
| | Urobilinogen | NEGATIVE / NORMAL |
| | | 1 - 2 - 2 - 2 - 2 |
| Roche Cobas U411 / | Bilirubin | NEGATIVE |
| Urisys I 100/1800/Miditron | | N. F. O. A. T. D. /F. |
| | Blood | NEGATIVE |
| | Ketones | NEGATIVE |
| | Leukocytes | NEGATIVE |
| | Nitrite | NEGATIVE |
| | pН | <5 – 6 |
| | Protein | NEGATIVE |
| | Specific Gravity | 1.015-1.025 RATIO |
| | Urobilinogen | NEGATIVE / NORMAL |
| | , , | |
| Roche Combur | Bilirubin | NEGATIVE |
| | Blood | NEGATIVE |
| | Glucose | NEGATIVE / NORMAL |
| | Ketones | NEGATIVE |
| | Leukocytes | NEGATIVE |
| | Nitrite | NEGATIVE |
| | pН | <5 – 6 |
| | Protein | NEGATIVE |
| | Specific Gravity | 1.015-1.025 RATIO |
| | Urobilinogen | NORMAL |
| | | |
| Siemens Clinitek Novus | Bilirubin | NEGATIVE |
| | Blood | NEGATIVE |
| | Glucose | NEGATIVE |
| | Ketones | NEGATIVE |
| | Leukocytes | NEGATIVE |
| | Nitrite | NEGATIVE |
| | pН | <5 – 6 |
| | Protein | NEGATIVE |
| | Specific Gravity | 1.02 - >1.03 RATIO |
| | Urobilinogen | 3.2 μmol/l / 0.2 EU/dl / 0.2 mg/dl |
| 0 | | A LEGATINE |
| Siemens Clinitest hCG | hCG | NEGATIVE |
| S: M. Li. | B.0. 1. | NECATIVE |
| Siemens Multistix | Bilirubin | NEGATIVE |
| | Blood | NEGATIVE |
| | Glucose | NEGATIVE |
| | Ketones | NEGATIVE |
| | Leukocytes | NEGATIVE |
| | Nitrite | NEGATIVE <5 – 6 |
| | pH Protein | NEGATIVE |
| | Specific Gravity | I.02 - >I.03 RATIO |
| | Urobilinogen | |
| | Urodilinogen | 3.2 μmol/l / 0.2 EU/dl / 0.2 mg/dl |

| MANUFACTURER / METHOD | ANALYTE | RANGE |
|------------------------------|------------|----------|
| Sysmex Meditape UC 10S & 12S | Bilirubin | NEGATIVE |
| | Blood | NEGATIVE |
| | Glucose | NEGATIVE |
| | Ketones | NEGATIVE |
| | Leukocytes | NEGATIVE |
| | Nitrite | NEGATIVE |
| | pН | <5 – 6 |
| | Protein | NEGATIVE |





| | Specific Gravity | 1.02 – 1.03 RATIO |
|-----------------------------------|------------------|---------------------------------------|
| | Urobilinogen | NORMAL |
| | Ü | |
| Sysmex Meditape UC 9A & 11A | Bilirubin | NEGATIVE |
| · | Blood | NEGATIVE |
| | Glucose | NEGATIVE |
| | Ketones | NEGATIVE |
| | Leukocytes | NEGATIVE |
| | Nitrite | NEGATIVE |
| | рН | <5 – 6 |
| | Protein | NEGATIVE |
| | Specific Gravity | 1.02 - 1.03 RATIO |
| | Urobilinogen | NORMAL |
| | | |
| Urit Medical Electronic | Bilirubin | NEGATIVE |
| | Blood | NEGATIVE |
| | Glucose | NEGATIVE |
| | Ketones | NEGATIVE |
| | Leukocytes | NEGATIVE |
| | Nitrite | NEGATIVE |
| | pН | <5 - 6 |
| | Protein | NEGATIVE |
| | Specific Gravity | 1.02 - 1.03 RATIO |
| | Urobilinogen | NORMAL |
| | | |
| YD Diagnostics / Biosys / Genesis | Bilirubin | NEGATIVE |
| , | Blood | NEGATIVE |
| | Glucose | NEGATIVE |
| | Ketones | NEGATIVE |
| | Leukocytes | NEGATIVE |
| | Nitrite | NEGATIVE (0.00 – 0.04 mg/dl) |
| | рН | <5-6 |
| | Protein | NEGATIVE |
| | Specific Gravity | 1.02 - 1.03 RATIO |
| | Urobilinogen | NORMAL / +/- / 0.1 mg/dl / 1.7 µmol/l |

Randox Teoranta, Meenmore,
Dungloe, Donegal,
F94 TV06, Ireland

11 May '23 me