

LIPID CONTROL - LEVEL 2 (LPD CONTROL 2)

Cat. No. LE2662 **Lot No.** 2256CH **Size:** 5 x 3 ml **Expiry:** 2017-03

INTENDED USE

This product is intended for *in vitro* use in the quality control of Direct HDL, Direct LDL, Lipoprotein (a), Apolipoprotein A-I, Apolipoprotein B, Cholesterol and Triglyceride methods on clinical chemistry systems.

SAFETY PRECAUTIONS AND WARNINGS

Human source material, from which this product has been derived, has been tested at donor level for the Human Immunodeficiency Virus (HIVI & HIV2) antibody, Hepatitis B surface antigen (HbsAg) and the Hepatitis C virus (HCV) antibody and found to be NON-REACTIVE. FDA approved methods have been used to conduct these tests. However, since no method can offer complete assurance as to the absence of infectious agents, this material and all patient samples should be handled as though capable of transmitting disease. For *in vitro* diagnostic use only.

STORAGE AND STABILITY

Unopened Lipid Control is stable until the expiry date printed on the product label, when stored between +2°C and +8°C. Once reconstituted, the components of the serum are stable for 7 days at +2°C to +8°C, and 4 weeks at -20°C when frozen once. The following exceptions apply: LP(a) is stable for 16 weeks at -20°C when frozen once. Values may drop by up to 10% for Direct LDL Cholesterol when stored for 4 weeks at -20°C.

PREPARATION FOR USE

Open the vial carefully, avoiding any loss of the material and reconstitute with 3 ml of distilled water. Replace the rubber stopper, close the vial and leave to stand for 30 minutes before use. Ensure that all traces of dry material are dissolved by swirling gently.

MATERIALS PROVIDED

Lipid Control - Level 2 5 x 3 ml

MATERIALS REQUIRED BUT NOT PROVIDED

Distilled water Volumetric pipette

VALUE ASSIGNMENT

Each batch of Lipid Control is submitted to a number of external laboratories. Values are assigned from a consensus of results obtained by these laboratories and internal testing conducted at Randox Laboratories Ltd.

If a method is unavailable, contact Randox Laboratories - Technical Services, Northern Ireland, tel: +44 (0) 28 9445 1070 or email Technical.Services@randox.com

10 Oct '14 ne



Cat. No. LE2662					
		44			
Analyte	unit	target	low	high	methods
Apolipoprotein A-1	g/l	1.50	1.23 123	1.77 177	Immunoturbidimetric
	mg/dl g/l	150	1.19	1.71	Nanhalamatria
	-	1.45 145	1.19		Nephelometric
Apolipoprotein B	mg/dl g/l	1.15	0.94	171	Immunoturbidimetric
	mg/dl	115	94.3	136	mmanotarbiametric
	g/l	1.19	0.98	1.40	Nephelometric
	mg/dl	119	97.6	1.40	Nephelometric
Cholesterol HDL - Cholesterol	mmol/l	5.87	5.11	6.63	Cholesterol Oxidase
	mg/dl	227	197	257	Cholesterol Oxidase
	mmol/l	5.44	4.73	6.15	Siemens Dimension
	mg/dl	210	183	237	Siemens dimension
	mmol/l	1.35	1.15	1.55	Direct Clearance Method
	mg/dl	52.1	44.4	59.8	Direct Glearance Method
	mmol/l	1.07	0.75	1.39	Phosphotungstic acid pptn.
	mg/dl	41.3	28.9	53.7	Phospholarigatic acid pptili.
	mmol/l	1.25	1.06	1.44	Direct HDL Immunoseparation
	mg/dl	48.3	40.9	55.7	Direct FIDE illimunoseparation
	mmol/l	1.20	1.02	1.38	Direct HDL PEGME
	mg/dl	46.3	39.4	53.2	DIIECT FIDE PEGINE
	mmol/l	1.34	1.14	1.54	Direct HDL PPD
	mg/dl	51.7	44.0	59.4	DIRECTION FED
	mmol/l	1.20	1.02	1.38	Direct HDL Roche 3rd generation
	mg/dl	46.3	39.4	53.2	Direct FIDE Roche Sid generation
	mmol/l	1.45	1.23	1.67	HDL - Ultra
	mg/dl	56.0	47.5	64.5	FIDE - Oilla
LDL - Cholesterol	mmol/l	4.02	3.42	4.62	Direct Clearance Method
	mg/dl	155	132	178	Direct Clearance Method
	mmol/l	3.54	3.01	4.07	Calculated
	mg/dl	137	116	158	Calculated
	mmol/l	3.48	2.96	4.00	Selective detergent methods
	mg/dl	3. 4 6 134	2.96 114	4.00 154	Ociconive detergent methods
Lipoprotein (a) Triglycerides	mg/dl	19.6	15.7	23.5	Immunoturbidimetric
	nmol/l	45.2	36.2	54.2	IIIIII I Oldi Didilli etile
	mmol/l	2.33	1.96	2.70	Lipase/GPO-PAP no correction
		2.33	1.96	2.70	Lipase/GFO-FAF No Correction
	mg/dl		1.97		Lingso/GK LIV no correction
	mmol/l	2.34		2.71	Lipase/GK UV no correction
	mg/dl	207	174	240	